

AN ANALYSIS OF THE SUBSET GRAPH OF A NEAR-RING USING MATHEMATICAL EQUATIONS

Dr. Niketa Sharma*

Assistant Professor in Mathematics
Hindu Kanya Mahavidyalaya, Jind, Haryana, India

Email ID: nikitaakshit@gmail.com

Accepted: 06.09.2022

Published: 01.10.2022

Keywords: Subset Graph, Near-Ring.

Abstract

This section introduces the concept of a subset graph of a near-ring. There are three distinct parts to this. The first portion handles the necessary basic definitions for the following sections, while the second section houses the primary results. Third, we discuss several subgraphs of the subset graph of a near-ring that have been launched. In this section N always denotes a zero-symmetric right abelian near-ring, unless otherwise specified. Since the introduction of the concept of Cayley graphs from groups, the study of graphs derived from algebraic structures has been of great interest to mathematicians. The zero-divisor graph of rings has attracted the attention of both algebraists and graph theorists in recent years.

Give N a chance to be a privilege abelian near-ring which is likewise zero symmetric.

We start with the accompanying definition

Definition: Give us a chance to consider a near-ring N where $(N, +)$ is an abelian group. Too give F a chance to be the arrangement of all non-empty subsets of N . We characterize the subset graph $F_{\mathfrak{N}}$ as the graph with every one of the members of F as vertices and any two distinct vertices A, B are adjacent if and just if $A+B = \{a + b : a \in A, b \in B\}$ is a correct N -subset of N .

Example: Give us a chance to consider the near-ring $N = \{0, a, b\}$ under the operations characterized by the accompanying tables.

+	0	a	b
0	0	a	b
a	a	0	a
b	b	a	0

.	0	a	b
0	0	0	0
a	0	a	a
b	0	b	b

Paper Identification



*Corresponding Author

1. INTRODUCTION

1.1 Preliminaries

Here we can see that $N, \{0, a\}, \{0, b\}$ are correct N -subsets of N . The graph $F_{\mathfrak{N}}$ is given beneath




Figure 1: The subset graph $F_{\mathfrak{R}}$

1.2 Main Results

Give N a chance to be a right-near ring and we indicate cardinality of N by α , for example $|N| = \alpha$. α may be of infinite cardinality.

We start our area with the accompanying outcome.

Theorem 1. For any directly near-ring N with $|N| \leq 2$, the graph $F_{\mathfrak{R}}$ is constantly associated. **Proof:** In the event that $|N| = 1$, at that point the proof is clear.

Let $|N| = 2$ and $N = \{0, a\}$.

The vertices $\{0\}$ and N are constantly associated. The vertices $\{a\}$ and $\{0\}$ are never adjacent, since $\{a\} + \{0\} = \{a\}$ which can't be a correct N -subset of N . Presently $\{a\} + N = N$ and plainly this is a correct N -subset of N . Hence the graph $F_{\mathfrak{R}}$ is associated.

Remark 1 In theorem 5.4.2.1, we have considered the near-ring as a zero-symmetric near-ring. It is fascinating to take note of that in the event that N isn't zero-symmetric, at that point with a similar adjacency relation for $|N| = 2$, the graph $F_{\mathfrak{R}}$ is a complete graph K_3 with diameter 1 and girth 3, for example consider the near-ring $\mathbb{Z}_2 = \{0, 1\}$ under expansion modulo 2 and $'\cdot'$ on \mathbb{Z}_2 is characterized as $a \cdot b = a, \forall a, b \in \mathbb{Z}_2$.

For $|N| > 2$, the graph $F_{\mathfrak{R}}$ is never completely disconnected. Anyway for any near-ring N , it is an extreme occupation to check whether there is an isolated vertex or not in the graph $F_{\mathfrak{R}}$. In the accompanying theorem we build up a few conditions

under which the graph $F_{\mathfrak{R}}$ never contains an isolated vertex.

Theorem 2: For $|N| > 2$, the graph $F_{\mathfrak{R}}$ is constantly associated on the off chance that one of the accompanying holds:

1. For any appropriate subset A , $A + N = N$.
2. For any two subsets A, B , $A + B = N$.
3. Every subset of N is a correct N -subset of N .

Proof: 1. Let $A \subset N$. Now $A + N \subseteq N$ and $(A + N)N \subseteq N \subseteq N$. It is certain that $(A + N)$ is a correct N -subset of N just if $N \subseteq (A + N)$, for example $A + N = N$. In this manner N is adjacent to each other vertex and consequently the graph $F_{\mathfrak{R}}$ is associated.

2. Let A, B be any two legitimate subsets of N with the end goal that $A + B = N$. Presently $(A + B)N = N \subseteq N = A + B$ and consequently A and B are adjacent. On the off chance that either $A = N$ or $B = N$, at that point from above (I) unmistakably A and B are adjacent and accordingly $F_{\mathfrak{R}}$ is associated

3. Let each subset of N is a correct N -subset of N . At that point from lemma 5.4.1.1 we have that any two vertices A, B in $F_{\mathfrak{R}}$ is adjacent to one another. Along these lines the graph $F_{\mathfrak{R}}$ is associated.

Remark 2 In the theorem, on the off chance that N fulfills either condition (ii) or (iii), at that point the coming about graph $F_{\mathfrak{R}}$ will be a complete graph $K_{2^{\alpha-1}}$. However If N fulfills condition (i) in theorem 5.4.2.2, at that point the graph $F_{\mathfrak{R}}$ contains a spanning subgraph isomorphic to the star graph $K_{1, 2^{\alpha-2}}$.

2. ON SOME SUBGRAPHS OF THE GRAPH $F_{\mathfrak{R}}$

In this area we examine some actuated subgraphs of the graph $F_{\mathfrak{R}}$. We likewise try to locate some graphical parameters like diameter, girth, chromatic number etc.

of these subgraphs.

I. Give us a chance to consider the subclass of the group of subsets of N which comprises of all the correct N -subsets of N . Give us a chance to signify this actuated subgraph of $F_{\mathfrak{R}}$ by $R_{\mathfrak{R}}$. We have the accompanying outcomes:

Theorem 1 The subgraph $R_{\mathfrak{R}}$ is a complete subgraph of $F_{\mathfrak{R}}$.

Proof: We realize that for any two right N -subsets of N state A, B , their entirety $A + B = \{a + b : a \in A \text{ and } b \in B\}$ is likewise a correct N -subset of N . Henceforth the announcement is clear.

Corollary 1 For any near-ring N we have $\text{diam}(R_{\mathfrak{R}}) \leq 1$.

Proof For any near-ring N , from theorem 5.4.3.1 we have, $R_{\mathfrak{R}}$ is a complete graph. Hence unmistakably $\text{diam}(R_{\mathfrak{R}}) \leq 1$

Conclusion 2: $gr(R_{\mathfrak{R}}) = 3 \text{ or } \infty$.

Proof: It is evident that the girth of a complete graph is either three or ∞ . Consequently $gr(R_{\mathfrak{R}}) = 3 \text{ or } \infty$.

II. For any near-ring N , an element $x \in N$ is said to be nilpotent if $x^t = 0$, for some $t \in \mathbb{Z}^+$. A subset S of N is known as a nilpotent subset of N if there exists a $k \in \mathbb{Z}^+$ with the end goal that $S^k = 0$ which implies $s_1 s_2 \dots s_k = 0$ for every $s_i \in S, i = 1, 2, 3, \dots, k$. A near-ring N is known as an emphatically semi-prime near-ring if N consider an incited subgraph of $F_{\mathfrak{R}}$ has no non-zero nilpotent invariant subset. Let us whose vertex set comprises of all the nilpotent subsets of N . Give us a chance to indicate this subgraph by $Nil_{\mathfrak{R}}$.

Lemma 1 An emphatically semi-prime near-ring N has no non-zero nilpotent left(right) N -subsets of N .

Theorem 2 On the off chance that N is unequivocally semi-prime, at that point the subgraph $Nil_{\mathfrak{R}}$ is the disjoint association of K_2 and K_1 's.

Proof. Let N be an emphatically semi-prime near-ring. By lemma 5.4.3.1., unmistakably N has no non-zero nilpotent right N -subsets of N . Therefore the subgraph contains a line $\{0\} - N$ and different vertices (assuming any) are isolated. Thus the outcome

Remark 1 A near-ring N is called regular if $\forall n \in N$, there exists $x \in N$ with the end goal that $nxn = n$. A near-ring N is called feebly regular if for any perfect i of N , each left i -subgroup A of $i, A_2 = A$. It can be seen that on the off chance that N is a pitifully regular

near-ring, at that point also theorem 5.4.3.2 holds.

As all regular near-rings are feebly regular so we can finally state that the subgraph $Nil_{\mathfrak{R}}$ is disjoint union of K_2 and K_1 's for a regular near-ring N .

3. LINE GRAPH ASSOCIATED TO GRAPH OF A NEAR-RING WITH RESPECT TO AN IDEAL

In this section we talk about line graph related to graph of a near-ring as for a perfect. This part comprises of two sections. The principal section manages the preliminary definitions required for the ensuing section and the second section contains the main results.

All through this section N indicates a directly near-ring except if generally expressed.

3.1 Preliminaries

Give N a chance to be a directly near-ring. We start with the accompanying definitions.

Definition 1: The near-ring N is called necessary on the off chance that it has no nonzero zero-divisors. N is called straightforward if its standards are $\{0\}$ and N

Definition 2: A perfect I of N is called prime, if for beliefs A, B of $N, AB \subseteq I$ infers either $A \subseteq I$ or $B \subseteq I$.

A perfect I is called semiprime if for any perfect J of $N, J^2 \subseteq I$ suggests that $J \subseteq I$.

Definition 3: A perfect I is called 3-prime if for $a, b \in N$ and $ab \subseteq I$ either $a \in I$ or $b \in I$. N is called 3-prime near-ring if $\{0\}$ is a 3-prime perfect of N .

Definition 4: A graph is said to be a cycle graph in the event that it comprises of a solitary cycle.

Definition 5: Give I a chance to be a perfect of N . The graph of N as for I is a graph with $V(N)$ as vertex set, and any two distinct vertices x and y are adjacent if and just if $xNy \subseteq I$ or $yNx \subseteq I$. This graph is indicated by $G_I(N)$.

Definition 6: Line graph of $G_I(N)$ is a graph with each edge of $G_I(N)$ as a vertex and any two distinct vertices are adjacent if and just if their comparing edges share a typical vertex in the graph $G_I(N)$. We indicate this line graph by $L(G_I(N))$. On the off chance that x, y be any two distinct vertices adjacent in the graph $G_I(N)$, at

that point the relating vertex in the line graph $L(G_I(N))$ is indicated by $[x, y]$.

Example 1: Let us consider \mathbb{Z}_4 the ring of integers modulo 4. The beliefs of \mathbb{Z}_4 are $I = \{0\}, J = \{0, 2\}$ and $K = \mathbb{Z}_4$. The graphs of \mathbb{Z}_4 regarding standards I, J and K and their comparing line graphs are appeared in figure 5.17 and figure 5.18 separately.




Figure 2: The graphs of $N = \mathbb{Z}_4$ with respect to ideals I, J and K




Figure 3: The Line graphs of $N = \mathbb{Z}_4$ with respect to ideals I, J and K

Example 2: Let us consider a near-ring $N = \{0, a, b, c\}$ under the two binary operations $+$ and $-$ characterized in the accompanying tables:

$+$	0	a	b	c
0	0	a	b	c
a	a	0	c	b
b	b	c	0	a
c	c	b	a	0

.	0	a	b	c
0	0	0	0	0
a	0	a	0	a
b	b	b	b	b
c	b	c	b	c

Here the beliefs of N are $I = \{0\}, J = \{0, b\}, K = \{0, b\}$ and $P = N$. The graphs of N as for these standards and their relating line graphs are appeared in figure 5.19.

3.2 Main Results

In this section we present some main results. Let N be a right-near ring with $|N| = n$ nowhere $|N|$ indicates the cardinality of N, n might be of infinite cardinality too.

We start our section with the accompanying outcome.

Remark 1 In the graph $G_I(N)$, 0 is constantly adjacent to the various vertices coming about somewhere around one edge and accordingly its comparing line graph will contain something like one vertex. Henceforth $L(G_I(N))$ can never be an empty graph.

Theorem 1 For any near-ring N , the graph $L(G_I(N))$ is constantly associated and $diam L(G_I(N)) \leq 3$.

Proof. In $G_I(N)$ the vertex 0 is adjacent to every vertex, so $G_I(N)$ is constantly associated and henceforth its line graph $L(G_I(N))$ is dependably an associated graph.

Give us now a chance to demonstrate the second part. Let $[x, y], [z, w]$ be any two vertices in $L(G_I(N))$.

At that point we can generally develop a way of length 3 as $[x, y] - [y, 0] - [0, z] - [z, w]$. In this manner $diam L(G_I(N)) \leq 3$

Remark 2 If N is straightforward and fundamental; at that point it is obvious from theorem 5.5.1.1. That the line graph $L(G_I(N))$ is either a regular graph of degree $2n - 4$ or a complete graph $K_n - 1$

The accompanying results give the girth of $L(G_I(N))$ under different conditions.

Theorem 2 For any near ring N let $I = \{0\}$. At that point $gr(L(G_I(N))) = \infty$ if and just if $N \cong \mathbb{Z}_2$ or $N \cong \mathbb{Z}_3$.

Proof: Let $N \cong \mathbb{Z}_2$ At that point $G_{\{0\}}(N)$ is an edge $0 - 1$. In this manner $(L(G_I(N)))$ contains just a single vertex and in this way $gr(L(G_I(N))) = \infty$. Again let $N \cong \mathbb{Z}_3$ length $2, 1 - 0 - 2$ thus $gr(L(G_I(N))) = \infty$.

On the other hand let $gr(L(G_I(N))) = \infty$. On the off chance that conceivable let N contains multiple elements. Assume $|N| = 4$. If $I = \{0\}$, at that point in $G_I(N)$ we will get an incited subgraph which is a star graph $K_{1,3}$. Thus the comparing line graph will contain a 4-cycle, which is an inconsistency. Thus $N \cong \mathbb{Z}_2$ or $N \cong \mathbb{Z}_3$.

Theorem 3: For any near-ring N whenever $G_I(N)$ contains a cycle, at that point $gr(L(G_I(N))) = 3$,

Proof: If $G_I(N)$ contains a 3 cycle, at that point we are finished.

Let $G_I(N)$ contains a 4 cycle. Consequently N contains atleast 4 elements. Since 0 is adjacent to each other vertex in $G_I(N)$, so there will be a star graph $K_{1,3}$ with focus vertex 0. Subsequently the comparing line graph will contain a subgraph isomorphic to K_3 . In this manner $(L(G_I(N)))$ will contain a 3-cycle thus $gr(L(G_I(N))) = 3$.

Next let $G_I(N)$ contains a 5-cycle. Then by the comparative contentions given above $G_I(N)$ contains a star graph isomorphic to $K_{1,4}$. Thus in the comparing line graph we will get a subgraph isomorphic to K_4 . Accordingly $gr(L(G_I(N))) = 3$

By The comparative contentions whenever $G_I(N)$ contains a n cycle, n is any non zero positive integer, in $(L(G_I(N)))$ there will be a complete subgraph isomorphic to K_{n-1} . Thus $(L(G_I(N)))$ contains a 3 cycle. Consequently $gr(L(G_I(N))) = 3$ accordingly the theorem is demonstrated.

Remark 3 The opposite of the theorem 5.5.2.3, does not hold always. For example the line graph $(L(G_I(N)))$ in figure 5.5.1.2. has girth 3, however its comparing graph $G_I(N)$ in figure 7.1.1 does not contain any cycle.

Theorem 4 The line graph $(L(G_I(N)))$ is a cycle graph if and just if either $NI = \mathbb{Z}_3$ or $N \cong \mathbb{Z}_4$ and $I \setminus \{0\}$.

Proof: Suppose $N = I \cong \mathbb{Z}_3$. It is obvious from the definition that $G_I(N)$ is a cycle of length 3 and henceforth the comparing line graph is a cycle graph

that is C_3 . Next assume $N = \mathbb{Z}_4$ and $I = \{0\}$. For this situation additionally $G_I(N)$ is a complete bipartite graph $K_{1,3}$ and it suggests that the relating line graph is a cycle of length 3

4. CONCLUSION

Near-rings are one of the generalized structures of rings. The study and research on near-rings is very systematic and continuous. Near-ring newsletters containing complete and updated bibliography on the subject are published periodically by a team of mathematicians (Editors: Yuen Fong, Alan Oswald, Gunter Pilz and K. C. Smith) with financial assistance from the National Cheng Kung University, Taiwan. These newsletters give an overall picture of the research carried out and the recent advancements and new concepts in the field. Conferences devoted solely to near-rings are held once every two years. There are about half a dozen books on near-rings apart from the conference proceedings. Above all there is an online searchable database and bibliography on near-rings. As a result the author feels it is very essential to have a book on Smarandache near-rings where the Smarandache analogues of the near-ring concepts are developed. The reader is expected to have a good background both in algebra and in near-rings; for, several results are to be proved by the reader as an exercise.

Bibliography

1. H.H.Abbass, M.N.Obaid, On A Primary Ideal and Semi Primary Ideal of a Near Ring, Journal of Kerbala University, 14 (2), pp. 163-170, 2016.
2. N. V. Nagendram, T. V. Pradeep Kumar and Y. V. Reddy, "IFP Ideal in Noetherian δ -Regular Near Rings" Int. J. Contemporar Math Vol. 2, No. 1, 2011.
3. Dheena and G.Satheesh Kumar, "Completely 2-primal Ideal in near ring", India, 2007.
4. G. L. Booth and K. Mogae, "Equiprime Ideal near-rings" M.Sc thesis University of the Nelson Mandela Metropolitan December 2008.

5. A. K.Jabbar and C. A. Ahmed, “On Almost Primary Ideals” International Journal of Algebra, Vol. 5, No. 13,pp: 627 - 636, 2011.
6. G. L. Booth and K. Mogae , “Equiprime Ideal near-rings” M.Sc thesis University of the Nelson Mandela Metropolitan December 2008.

